



Legal status:

Dietetic supplement (in accordance with 767/2009/EG and RL 2008/38/EG), delivery to the final consumer

Additives per liter:

Vitamin A (3a672b)	50 000 000	I.U.
Vitamin D3 (3a671)	5 000 000	I.U.
Vitamin E (3a700)	50 000	mg
Sodium ascorbate (1b301)	100 000	mg
(Vitamin C)		-

Density:

Approx. 1.07 g/cm³

Solubility: Completely soluble in water

Animal categories:

For poultry

Recommended dosage:

100 ml per 1000 lifters of drinking water

The opinion of a nutrition expert or veterinarian should be sought before using the product or before extending its period of use

The daily total intake of vitamin A and vitamin D3 must be observed. The maximum content of vitamin A per kg of complete feed is 20,000 I.U. for chickens < 14 days and for turkeys < 28 days; for other poultry maximum content of vitamin A is 10,000 I.U./kg complete feed. The maximum content of vitamin D3 per kg of complete feed is 5000 I.U. for broilers and 3000 I.U. per kg for other poultry.

Particular nutritional purpose:

Compensation of vitamin A and vitamin D3 deficiency in impaired absorption (malabsorption) and thus avoidance of disorders of bone and shell quality. To be used during the first two weeks after hatching.

Support preparation for oestrus and reproduction, in female animals during oestrus and in male animals during reproductive activity.

Recommended use:

Shake well before application, do not use in pure form. Dissolve 100 ml in min. 10 litres of water and fill up to the required quantity.

Storage:

Keep tightly sealed and protected from direct sunlight in a cool (max. 25°C) and dry place

Shelf life:

Guaranteed contents for 24 months if kept under proper storage conditions in original packaging

Packaging:

Bottle of 1 litre in carton 12 x 1 litres Canister of 5 litres in carton 4 x 5 litres

Code number:

47721 (1 litre) 47722 (5 litres)

Kaesler Nutrition

Kaesler Nutrition GmbH Zeppelinstraße 3 27472 Cuxhaven, Germany Phone: +49 (0)4721 5904-0